

Asteroseismology of the mild Am δ Sct star HD 118660: *TESS* photometry and modelling

Mrinmoy Sarkar¹,^{1,2} Santosh Joshi,¹ Marc-Antoine Dupret,³ Otto Trust⁴, Peter De Cat⁵, Eugene Semenko⁶, Patricia Lampens,⁵ Aruna Goswami,⁷ David Mkrtychian⁶, Drisya Karinkuzhi,⁸ Ilya Yakunin⁹ and Archana Gupta²

¹Aryabhata Research Institute of Observational Sciences, Manora Peak, Nainital 263002, India

²Department of Applied Physics, M.J.P. Rohilkhand University, Bareilly, Uttar Pradesh 243006, India

³Space Sciences, Technologies and Astrophysics Research (STAR) Institute, University of Liège, B-4000 Sart Tilman, Belgium

⁴Department of Physics, Mbarara University of Science and Technology, P.O. Box 1410, Mbarara, Uganda

⁵Royal Observatory of Belgium, Ringlaan 3, B-1180 Brussels, Belgium

⁶National Astronomical Research Institute of Thailand, Mae Rim, Chiang Mai-50180, Thailand

⁷Indian Institute of Astrophysics, Koramangala, Bangalore 560034, India

⁸Department of Physics, University of Calicut, Thenhipalam, Malappuram 673635, India

⁹Special Astrophysical Observatory, Russian Academy of Sciences, Nizhniy Arkhyz 369167, Russia

Accepted 2024 September 25. Received 2024 September 2; in original form 2023 September 29

ABSTRACT

We present the results of an asteroseismic study of HD 118660 (TIC 171729860), being a chemically peculiar (mild Am) star exhibiting δ Scuti (δ Sct) pulsations. It is based on the analysis of two sectors of time-series photometry from the space mission *TESS* (*Transiting Exoplanet Survey Satellite*) and seismic modelling. It yielded the detection of 15 and 16 frequencies for *TESS* sectors 23 and 50, respectively. The identified pulsation modes include four radial ($\ell = 0$) and five dipolar ($\ell = 1$) ones. The radial modes are overtones with order n ranging from 3 and 6. Such high values of n are theoretically not expected for stars with the effective temperature of HD 118660 ($T_{\text{eff}} \approx 7550$ K) located near the red edge of the δ Sct instability strip. To estimate the asteroseismic parameters, we have generated a grid of stellar models assuming a solar metallicity ($Z = 0.014$) and different values for the convective overshooting parameter ($0.1 \leq \alpha_{\text{ov}} \leq 0.3$). We conclude that the analysis of the radial modes is insufficient to constrain α_{ov} and Z for δ Sct stars. The value for the equatorial velocity of HD 118660 derived from the seismic radius and the rotational frequency is consistent with values found in the literature.

Key words: asteroseismology – convection – diffusion – stars: activity – stars: chemically peculiar – stars: evolution.

1 INTRODUCTION

Asteroseismology uses stellar pulsations, which appear as periodic variations of brightness, radial velocities, and/or line profiles, to probe the internal structure and evolution of stars. Pulsating stars are ubiquitous in the Hertzsprung–Russell (H–R) diagram (Kurtz 2022). The δ Scuti (δ Sct) stars are of particular interest for asteroseismology for several reasons. The richness of their pulsation spectra allows for the probing of several different internal layers. Their brightness makes them accessible for high-precision observations using both photometry and spectroscopy.

The masses of δ Sct variables typically range from 1.5 to 2.5 M_{\odot} (Bregier 1979; Goupil et al. 2005; Guzik 2021; Kurtz 2022). They are located in the H–R diagram where the classical instability strip intersects with the main sequence (Dupret et al. 2004, 2005), confined to the range of effective temperatures (T_{eff}) nearly between 7000 to 9000 K (Uytterhoeven et al. 2011). In this region, stellar

interiors change from radiative cores with thick convective envelopes ($M \lesssim 1 M_{\odot}$) to the convective cores with thin radiative envelopes ($M \gtrsim 2 M_{\odot}$). Hence, the study of δ Sct stars can give important insights into the details of this transition.

Thanks to recent space missions like *Kepler* (Christensen-Dalsgaard et al. 2008), *K2* (Howell et al. 2014), and the *Transiting Exoplanet Survey Satellite* (*TESS*; Ricker et al. 2015), we have a plethora of high-precision photometric data. Analysis of these light curves shows that the δ Sct stars mostly pulsate in pressure (p) modes with observed oscillation frequencies ranging from ~ 5 to 100 d^{-1} (Guzik, Garcia & Jackiewicz 2019; Murphy et al. 2019; Bedding et al. 2020). At the same time, the space-based photometric observations have revealed that a significant fraction of δ Sct stars simultaneously exhibit frequencies that are typical for γ Doradus (γ Dor) stars, making them hybrids (e.g. Balona 2014; Hey & Aerts 2024). The multiperiodic pulsational nature of δ Sct stars make them an ideal test bed for asteroseismology.

The δ Sct stars pulsate in low-order radial and non-radial overtones. The corresponding observed amplitudes can be as low as a few tens of μ -mag (low-amplitude δ Sct stars; Murphy 2014) or as high as

* E-mail: mrinmoys235@gmail.com

a few tenths of a magnitude (high-amplitude δ Sct stars; Lee et al. 2008). The opacity mechanism operating in the He II ionization zone is believed to be the primary source for driving pulsations in δ Sct stars, although other mechanisms are likely to contribute (Houdek 2000; Antoci et al. 2014; Murphy et al. 2020). Stellar metallicity is one of the key drivers in the excitation of their pulsations (Guzik, Fryer & Fontes 2018).

The influence of anomalous chemical abundances on stellar pulsations is not yet fully understood. Pulsating A-type stars, including the δ Sct variables, share the same region of the main sequence with objects whose key feature is an abnormal chemical composition. These so-called chemically peculiar (CP) stars differ from their chemically normal counterparts as their spectra exhibit strengthened or weakened lines of specific elements (Preston 1974). The CP stars constitute about 10 per cent–15 per cent of the population of stars with the same spectral type (Renson & Manfroid 2009).

The diffusion of atoms in stellar atmospheres is considered to be responsible for the formation of peculiarities in CP stars (Michaud 1970; Michaud, Alecian & Richer 2015). Some elements, particularly He, N, and O, settle in the atmosphere into the layers below the surface while others, such as Mn, Sr, Y, and Zr, are levitated out to the surface (Deal & Monier 2020). When various mixing effects like convection, rotation, turbulence, or mass loss, are absent or do not play an important role, atomic diffusion can develop into a rather stable configuration with a non-uniform horizontal or vertical distribution of atomic species. A stable magnetic field can also provide the required stability. As a result, from an observational point of view, such a CP star may appear to have an abnormal chemical composition and, hence, metallicity. In reality, its average composition may not substantially differ from a non-peculiar star of the same origin and age (Murphy et al. 2020).

The CP stars can be divided into smaller subgroups based on their most prominent chemical anomalies, binarity status, and magnetic properties. In this region of the H-R diagram, a star is classified as a magnetic CP star if it possesses a strong, stable, and globally organized magnetic field with a fossil origin, simple structure and surface strength of at least ≈ 300 G. Amongst the magnetic CP stars, a small fraction pulsates. They are known as rapidly oscillating Ap (roAp) stars because of their short pulsation periods ranging between approximately 5 and 30 min (Kurtz 1982). Only about 100 of them are currently known (Holdsworth et al. 2021, 2024, and references therein). The metallic-line (Am) peculiar stars are more numerous and are characterized by an underabundance of light elements (He, Ca, and/or Sc) in combination with an overabundance of iron-group metals. Like δ Sct stars, Am stars are generally considered to be non-magnetic CP stars (Fossati et al. 2007, 2008; Donati & Landstreet 2009). However, a limited number of studies have reported on the detection of a magnetic field in δ Sct and Am stars (e.g. Kurtz et al. 2008; Neiner & Lampens 2015; Hubrig et al. 2023).

Axial rotation is an important factor governing stellar pulsations, but it is not straightforward to fully take it into account when analysing observations and calculating theoretical models. In the case of a slowly rotating pulsating star, the axial rotation already causes a shift in the observed frequencies (Ledoux 1949; Hansen, Cox & van Horn 1977). This scenario becomes more complicated when fast rotation also alters the shape of the star into an oblate spheroid, as it changes the stellar interior and potentially influences other physical processes that contribute to the pulsation mechanism. Convection is one of the physical processes that also affect pulsation. When convective core overshooting occurs, the core boundary is extended, leading to extra mixing in the stellar interior and enhancing the main-sequence lifetime of a star as the core gets more fuel. There are very

few studies about the impact of rotation on the corresponding overshooting parameter α_{ov} . Therefore, in-depth asteroseismic studies of rapidly rotating stars are required to test and improve our current understanding of the driving of pulsations. In contrast to the magnetic roAp stars, their non-magnetic pulsating counterparts usually show much weaker modulation of the brightness, which is observed in the most precise space-based light curves and can be attributed to the axial rotation (e.g. Balona 2011, 2019). Hybrid δ Sct- γ Dor pulsators and unresolved binaries can show peaks in the same frequency region, where the rotationally induced modulation is expected (e.g. Sikora et al. 2019; Rodríguez et al. 2020; Sikora, Wade & Rowe 2020). This limits the selection of rapidly rotating stars to measure the broadening of absorption lines in spectra or search for rotationally split frequencies in time-series data. The spectroscopic survey of Solano & Fernley (1997) revealed that δ Sct stars show a broad distribution in their projected rotational velocity ($v \sin i$) and they found an average value $\langle v \sin i \rangle$ of 97 km s^{-1} for the low-amplitude pulsators and a maximum value exceeding 200 km s^{-1} .

More than two decades ago, the Nainital-Cape Survey was initiated by astronomers from India and South Africa to detect and study the pulsational variability in Ap and Am/Fm stars (e.g. Ashoka et al. 2000; Martinez et al. 2001; Joshi et al. 2003, 2006, 2009, 2010, 2012, 2016, 2017). Meanwhile, the project developed into a multinational collaborative venture for which the most recent results have been published by Trust et al. (2020, 2021, 2023) and Joshi et al. (2022). HD 118660 is one of the targets of this survey for which photometric observations were gathered that were analysed by Joshi et al. (2006). They discovered δ Sct-type pulsations with prominent pulsation periods of ~ 1 and 2.52 h. The subsequent combined spectroscopic and spectropolarimetric study of Joshi et al. (2017) revealed that this star has a subsolar Ca abundance of -5.92 ± 0.32 and shows an excess in Fe of -4.76 ± 0.23 , where the solar values are -5.73 and -4.53 , respectively and elements of the lanthanides group typical for the mild metallic-line stars. Given the absence of a detectable magnetic field in HD 118660 and derived abundances, this object deserves special attention due to exceptionally high $v \sin i$ -value of $108 \pm 8 \text{ km s}^{-1}$ close to the upper limit ($\sim 120 \text{ km s}^{-1}$) observed for the Am stars (Abt 2009). Moreover, these authors also observed line profile variations in two consecutive short-exposure spectra with a high signal-to-noise ratio (SNR). These facts motivated us to perform a comprehensive study of HD 118660 using modern observational techniques and up-to-date theoretical models. In the meantime, HD 118660 has been subjected to a multisite campaign to acquire time-resolved spectroscopic data. The results based on these data will be published in a forthcoming paper (Semenko et al., in preparation). A preliminary analysis of the *TESS* time-series data for the two currently available sectors shows a rich frequency spectrum with clear evidence of amplitude modulation due to the beating of two close frequencies (Sarkar, Joshi & de Cat 2024). In this paper, we perform a comprehensive analysis of these *TESS* light curves and theoretical modelling in which we investigate the role of core overshooting in the seismic activity of this star.

The structure of the paper is as follows. Section 2 describes the observational data and processing. A detailed presentation of the asteroseismic modelling and results are given in Section 3. In Section 4, we end with an overview of the main conclusions drawn from our study and a discussion of their relevance.

2 OBSERVATIONS AND DATA REDUCTION

TESS (Ricker et al. 2015) is a NASA space-based observatory launched on 2018 April 18 into a 13.7-d cislunar orbit maintained by a

2:1 lunar resonance. The primary goals of the mission are finding new exoplanets and characterizing previously known systems using the transiting method. Moreover, the data collected by *TESS* are ideally suited for detailed asteroseismic studies of pulsating variables.

HD 118660 was observed by the *TESS* in sectors 23 (S23; from 2020 March 19 to April 15) and 50 (S50; from 2022 March 26 to April 22).

We used the PYTHON module LIGHTKURVE (Lightkurve Collaboration 2018) to retrieve the *TESS* data of HD 118660 from the Barbara A. Mikulski Archive for Space Telescopes (MAST; Fleming 2019). This object was observed in short cadence mode with a time sampling of 2 min. The PDCSAP (Pre-Data Conditioned Simple Aperture Photometry) flux data, corrected for long-term trends and systematic errors, were considered for analysis. The resulting light curves and the corresponding frequency spectra of HD 118660 generated using Discrete Fourier Transform (DFT) in PERIOD04 (Lenz & Breger 2005) are depicted in the upper and lower panels of Fig. 1 for sectors 23 and 50, respectively.

The variations corresponding to the significant frequencies were extracted from the *TESS* light curves. We used the software period04 for this purpose, which performs DFT of the input time-series data. The significance evaluation of frequency peaks is based on an SNR criterion. As recommended by Baran, Koen & Pokrzywka (2015), only frequencies with an amplitude exceeding five times the mean noise level ($\text{SNR} > 5$) were retained, where the mean noise level is evaluated near the Nyquist limit of the frequency spectra.

The uncertainties of the frequencies are calculated as

$$\sigma_f = 0.44 \frac{1}{T_{\text{obs}}} \frac{1}{\text{SNR}}, \quad (1)$$

where T_{obs} is the time span of the observations (e.g. Montgomery & O'Donoghue 1999; Kjeldsen & Bedding 2012). To determine the errors of the amplitudes and phases, we have performed a 10000-step Monte Carlo test in PERIOD04. The resulting significant frequencies detected along with their corresponding amplitudes, phases, and SNR values are tabulated in Table 1. In our analysis, we considered frequencies larger than 1 d^{-1} for both S23 and S50 observations. As for S23, there have no significant frequencies below that level and for S50 the noise level is too high.

3 ASTEROSEISMIC MODELLING

3.1 Modelling parameters

In our analysis, we performed model-based seismic analysis, estimating the seismic parameters followed by mode identification of the pulsation frequencies. The stellar evolution models were created using the Code Liégeois d'Evolution Stellaire (CLÉS; Scuflaire et al. 2008a). The step-wise radial eigenfrequencies were calculated using the Liège Oscillation Code (Scuflaire et al. 2008b). The grid of models encompasses a mass range of $1.10\text{--}2.00 M_{\odot}$ with a step size of $0.05 M_{\odot}$. The metallicity (Z) varies from 0.002 to 0.020 with a step of 0.002. Finally, the grid incorporates a convective overshooting parameter (α_{ov}) values of 0.1, 0.2, and 0.3. The stellar model calculation was started with an initial H-abundance of 75 per cent assuming that the star is on the zero-age main sequence (ZAMS) and continued until the star reaches the red giant branch. The estimated rotation velocity of the star is nearly 28 per cent of the critical value. For δ Sct stars a 2D numerical approach shows that that rotation velocity below the 50 per cent of the critical value, the low-order modes ($\ell = 0, 1$) normalized by the large separation, these

modes remain fairly constant (Reese 2022). Therefore, in our model calculations, we have not considered the effect of rotation.

3.2 Mode identification

The δ Sct variables pulsate in low-order overtones of p modes with pulsation frequencies above 5 d^{-1} . Two techniques were used to identify the modes: the interpretation of the Échelle diagram and the analysis of frequency ratios. The analysis of the frequency ratios have done by theoretical Petersen (1973, 1978) diagram. The following subsections provide the details of the applied procedures.

3.2.1 Échelle diagram

In the asymptotic regime, where $n \gg l$, the modes of a given ℓ are approximately equally spaced in the frequency domain by the value of parameter $\Delta\nu$, which is known as large frequency separation. The inverse value of $\Delta\nu$ is the time taken by a sound wave to traverse the star. The frequencies ν are related to the order n and degree ℓ of the pulsation modes and to the large frequency separation $\Delta\nu$ through the asymptotic equation of the form (Tassoul 1980)

$$\nu_{n,\ell} = \Delta\nu (n + \ell/2 + \epsilon), \quad (2)$$

where ϵ is a constant of order unity. It is also well established that the oscillation frequencies (and therefore the large separation $\Delta\nu$) scale as the square root of the mean stellar density ($\bar{\rho}$) by the empirical relation $\Delta\nu \propto \sqrt{\bar{\rho}}$ (Aerts, Christensen-Dalsgaard & Kurtz 2010).

Equation (2) demonstrates that the radial and non-radial modes appear repetitively with the same overtones for lower ℓ in the Fourier transform. However, the δ Sct stars do not pulsate in the asymptotic regime (Bedding et al. 2020). Furthermore, HD 118660 is a fast-rotating star and this fact seriously restricts the use of equation (2) for detailed seismic analysis. In our study, we refer to equation (2) only during the identification of radial modes and for the first estimation of the large frequency separation.

The Échelle diagram is an important tool for analysing spectra of stellar pulsations. In this diagram, the frequency spectrum is sliced into equal chunks with a width of $\Delta\nu$ and stacked vertically towards higher frequencies (Bedding et al. 2020). We used a dynamic Échelle interface (Hey & Ball 2022) to fine-tune the value of $\Delta\nu$. In Fig. 2, we show frequency spectra of HD 118660 in Échelle format. The best visually aligned pattern with two vertical ridges corresponding to the angular degrees $\ell = 0$ (left) and 1 (right) is achieved for $\Delta\nu = 4.31 \text{ d}^{-1}$.

For the frequencies detected in the *TESS* data of both sectors that are situated on the radial ridge of the Échelle diagram, we have introduced an additional identifier in the second column of Table 1. The frequencies aligned in the vertical direction correspond to the consecutive overtones. The estimated value of $\Delta\nu$ and the mean stellar density computed from the best-fitting model for HD 118660 agree well with the corresponding characteristics published in Bedding et al. (2020) for a group of δ Sct stars, as shown in Fig. 3.

3.2.2 Frequency ratios

For the frequency ratio analysis, we only considered the observed frequencies associated with radial modes ($\ell = 0$), being $f_{r,1}$, $f_{r,2}$, $f_{r,3}$, and $f_{r,4}$ (second column of Table 1). We calculated the frequency ratios for different pairs of radial orders under the assumption that the star is located on the main sequence. In Fig. 4, we show theoretical Petersen diagrams obtained for $Z = 0.014$ and $\alpha_{\text{ov}} = 0.1, 0.2, 0.3$




Figure 1. Top row: the left panel displays the *TESS* light curve observed in short cadence mode (2-min sampling) in sector 23, and its corresponding Fourier transform is given in the right panel. Bottom row: Idem, but for the data obtained in *TESS* sector 50.

Table 1. The significant frequencies detected in the variations of the 2-min cadence *TESS* data of sectors 23 (top part) and 50 (bottom part). Each record lists the identifier (ID), the frequency value f and its error σ_f , the degree ℓ and order n of the pulsation mode it has been identified with (if available), the amplitude A and its error σ_A , the phase ϕ and its error σ_ϕ , and the SNR value of the corresponding peak in the DFT. An additional ID is given in the second column for the radial modes detected in both sectors (IDrad).

ID	IDrad	f (d ⁻¹)	σ_f (d ⁻¹)	(ℓ , n)	A (mmag)	σ_A (mmag)	ϕ	σ_ϕ	SNR
Sector 23									
$f_{23,1}$	$f_{r,1}$	30.9202	0.0001	(0, 5)	0.5911	0.0033	0.4141	0.0008	131
$f_{23,2}$	$f_{r,2}$	26.6894	0.0002	(0, 4)	0.3691	0.0032	0.5203	0.0013	82
$f_{23,3}$		24.3829	0.0002	–	0.2794	0.0037	0.2817	0.0021	62
$f_{23,4}$		24.4431	0.0004	(1, 3)	0.2053	0.0037	0.1425	0.0028	46
$f_{23,5}$		28.6945	0.0005	(1, 4)	0.1561	0.0032	0.1425	0.0032	35
$f_{23,6}$		32.9514	0.0005	(1, 5)	0.1467	0.0032	0.7954	0.0035	33
$f_{23,7}$	$f_{r,3}$	22.1567	0.0007	(0, 3)	0.1101	0.0035	0.3288	0.0051	24
$f_{23,8}$		22.0904	0.0010	–	0.0805	0.0035	0.6042	0.0064	18
$f_{23,9}$		37.2650	0.0012	(1, 6)	0.0666	0.0032	0.4172	0.0077	15
$f_{23,10}$		34.4361	0.0014	–	0.0578	0.0032	0.4813	0.0090	13
$f_{23,11}$	$f_{r,4}$	35.0489	0.0016	(0, 6)	0.0479	0.0032	0.6645	0.0107	11
$f_{23,12}$		40.9176	0.0018	–	0.0431	0.0032	0.9283	0.0117	10
$f_{23,13}$		41.8077	0.0020	(1, 7)	0.0394	0.0033	0.2859	0.0134	9
$f_{23,14}$		41.8920	0.0020	–	0.0384	0.0014	0.1522	0.3292	9
$f_{23,15}$		1.0846	0.0030	–	0.0246	0.0121	0.4911	0.2345	6
Sector 50									
$f_{50,1}$	$f_{r,1}$	30.9204	0.0001	(0, 5)	0.6121	0.0047	0.1694	0.0012	107
$f_{50,2}$	$f_{r,2}$	26.6889	0.0002	(0, 4)	0.3741	0.0047	0.7713	0.0020	65
$f_{50,3}$		24.3792	0.0002	–	0.2806	0.0057	0.8311	0.0033	49
$f_{50,4}$		24.4383	0.0004	(1, 3)	0.2120	0.0058	0.2648	0.0041	33
$f_{50,5}$		28.6937	0.0005	(1, 4)	0.1586	0.0048	0.8905	0.0053	35
$f_{50,6}$		32.9518	0.0006	(1, 5)	0.1553	0.0047	0.2479	0.0048	27
$f_{50,7}$	$f_{r,3}$	22.1629	0.0007	(0, 3)	0.1048	0.0057	0.9433	0.0085	24
$f_{50,8}$		34.4345	0.0010	–	0.0664	0.0049	0.9410	0.0115	18
$f_{50,9}$		41.8914	0.0012	–	0.0669	0.0058	0.8558	0.0264	15
$f_{50,10}$		22.0905	0.0013	–	0.0858	0.0057	0.7898	0.0102	13
$f_{50,11}$		37.2663	0.0016	(1, 6)	0.0573	0.0047	0.8712	0.0131	11
$f_{50,12}$	$f_{r,4}$	35.0481	0.0018	(0, 6)	0.1169	0.0099	0.4763	0.0296	10
$f_{50,13}$		40.9137	0.0020	–	0.0491	0.0006	0.3884	0.0778	9
$f_{50,14}$		33.4568	0.0020	–	0.0438	0.0006	0.4374	0.0425	9
$f_{50,15}$		29.1510	0.0025	–	0.0312	0.0006	0.1043	0.1100	7
$f_{50,16}$		1.0983	0.0035	–	0.0239	0.0118	0.1981	0.1160	5




Figure 2. Left panel: the Échelle diagram depicts frequencies derived from the *TESS* light curve in S23. The circular points represent radial ($\ell = 0$) overtones, and the triangular points are dipolar overtones ($\ell = 1$). Right panel: Échelle diagram with same observed frequencies for S50.




Figure 3. A comparison of HD 118660 (asterisk) to a group of δ Sct stars (circles with errorbar) listed in Bedding et al. (2020) that indicates that the large separation $\Delta\nu = 4.31 \text{ d}^{-1}$ found in our study for HD 118660 follows the $\Delta\nu$ versus $\bar{\rho}$ relation within the 1σ spread. The error in $\bar{\rho}$ corresponds to the 1σ spread of the mean density of best-fitting models shown in Fig. 5.

(in red, blue, and green colours, respectively). We use the notation f_{n+1} to refer to the theoretical frequency of the n th overtone radial mode. Each panel of Fig. 4 shows the results for another radial mode in the numerator of the frequency ratios: the first overtone in the left panel ($n = 2$), the second overtone in the middle panel ($n = 3$), and the third overtone in the right panel ($n = 4$). The dots represent the frequency ratios of the observed radial modes, being

$f_{r,3}/f_{r,2} \sim 0.8303 \text{ d}^{-1}$ (upper dot), $f_{r,3}/f_{r,1} \sim 0.7167 \text{ d}^{-1}$ (middle dot), and $f_{r,3}/f_{r,4} \sim 0.6323 \text{ d}^{-1}$ (lower dot). The best visual match is found in the middle panel of Fig. 2, corresponding to the second overtone ($n = 3$) in the numerator.

We did the same exercise for the other values of the overshooting parameter α_{ov} . It causes small variations of the theoretical frequency ratios for frequencies below 15 d^{-1} , while the frequency ratio values in the flat part of the diagram remain unchanged. As the detected radial mode with the lowest frequency is located in the flat region of the Peterson diagrams, we conclude that the identification of the radial overtones does not depend on the value of α_{ov} .

Overtones of orders outside the plotted range are not included intentionally. The possibility of finding another correlation with the data remains, but we leave it unexplored as, in this case, the large separation must be an integer multiple or fraction of 4.31 d^{-1} .

The overall conclusion of our efforts to identify the modes (Sections 3.2.1 and 3.2.2) is that the radial overtones of orders $n = 3$ – 6 are present in the observed variations of the *TESS* data. This interpretation was added in Fig. 2. These radial overtones are almost equidistant in frequency.

3.3 Stellar models

Our study is based upon a grid of 570 evolution tracks calculated for different values of the independent parameters M (19 values), Z (10 values), and α_{ov} (3 values). We used the seismic- χ^2 distribution from Murphy et al. (2021) to find minima of the χ^2 values from each of the grid mentioned earlier.

In Fig. 5, the results of the seismic modelling are shown as the scattered plots of the mean stellar density M/R^3 versus mass M , radius R , and age t . The stellar models, corresponding to the




Figure 4. The evolution of frequency ratios for radial modes ($\ell = 0$) of consecutive overtones, plotted using theoretical models with $Z = 0.014$ and $\alpha_{ov} = 0.1, 0.2, 0.3$ (in different colors). The frequencies in the numerator correspond to the first overtone f_2 ($n = 2$; left panel), second overtone f_3 ($n = 3$; middle panel), and third overtone f_4 ($n = 4$; right panel). The dots denote the frequency ratios of the observed radial modes with $f_{r,3}$ in the numerator.




Figure 5. Illustration of the relationship between M/R^3 and the parameter mass (left panel), radius (middle panel), and age (right panel). The circular points of different colors represent the models limited by the criteria described in Section 3.3 for $\alpha_{ov} = 0.1, 0.2$, and 0.3 , respectively. The smallest dots represent all stellar models, while the medium and big sizes indicate models limited within 3σ and 1σ from the observed position in the H-R diagram, respectively.

convective overshooting parameters $\alpha_{ov} = 0.1, 0.2, 0.3$, are plotted in red, blue, and green, respectively. The models constrained within 3σ and 1σ from the observed position in the H-R diagram are marked using the medium and big markers. In the following subsections, we outline the details of the modelling.

3.3.1 Overshooting parameter

The overshooting parameter α_{ov} strongly depends upon the mass for the low- and intermediate-mass stars (Umezu 1995; Claret & Torres 2016, 2017, 2018, 2019). The value of α_{ov} increases almost linearly for stars in the mass range $M \sim 1.2\text{--}2.0 M_\odot$ and saturates near $\alpha_{ov} \sim 0.22$ for stars with M up to $4.4 M_\odot$. Convective overshooting plays a vital role in stellar evolution as it extends the lifetime of stars on the main sequence. The significance of overshooting increases with age and mass while the exact value of α_{ov} remains not well established (Lovekin & Guzik 2017; Claret & Torres 2018; Johnston 2021; Dornan & Lovekin 2022). Indeed, the uncertainty of α_{ov} at the saturation point can reach up to 0.4 (Constantino & Baraffe 2018). In light of these observations and considering the spectral type of

HD 118660, our choice to constrain α_{ov} between 0.1–0.3 for the seismic analysis appears to be well justified.

The overshooting parameter α_{ov} was implemented in the stellar evolution code CLÉS to calculate the size of the overshooting zone r_{ov} , given by

$$r_{ov} = r_c \pm \alpha_{ov} \min(H_p, h), \quad (3)$$

where r_c is the radius of the convective zone at the boundary, h is its size, and H_p is pressure scale height.

To demonstrate the importance of α_{ov} as a seismic parameter, we show two different sets of evolution tracks computed for $\alpha_{ov} = 0.1$ (solid line) and 0.3 (dotted line) for Z fixed at 0.014 in the right panel of Fig. 6. The position of HD 118660 is shown with black mark. One can notice that the overshooting parameter has an increasing effect on the seismic age in the main-sequence phase. With higher values of α_{ov} , the more intensive conveyance of fresh hydrogen in the stellar core extends the stellar lifespan. As a result, the seismic age increases with the overshooting parameter (Table 2). However, the impact of α_{ov} on the rest of seismically estimated stellar parameters is less




Figure 6. Left panel: the evolution tracks corresponding to the global metallicities $Z = 0.008$ (dotted lines) and $Z = 0.014$ (solid lines) computed for $\alpha_{ov} = 0.2$. Right panel: the evolution tracks calculated for $\alpha_{ov} = 0.1$ (solid lines) and $\alpha_{ov} = 0.3$ (dotted lines) for solar metallicity ($Z = 0.014$). The black error bar represents the position of HD 118660 on the H-R diagram.

Table 2. The final values of metallicity, mass, radius, and age calculated in the current study using the mean values of models constrained within 1σ error of the observed $\log(L/L_\odot)$ and $\log T_{\text{eff}}$. And the error corresponding to the mass, age, and v_{eq} mentioned here for stellar models with α_{ov} showing largest spread.

α_{ov}	$\frac{M}{M_\odot}$	$\frac{R}{R_\odot}$	Age (Gyr)	$\frac{M}{R^3}$ [solar]	v_{eq} (km s $^{-1}$)
0.1	1.77 ± 0.04	2.06 ± 0.01	1.15 ± 0.07	0.20	114 ± 1
0.2	1.80 ± 0.04	2.07 ± 0.01	1.21 ± 0.07	0.20	115 ± 1
0.3	1.75 ± 0.04	2.06 ± 0.01	1.35 ± 0.07	0.20	114 ± 1

significant. For example, our study shows that the seismic parameters of HD 118660, are independent of α_{ov} except for the stellar age.

3.3.2 Metallicity

The stellar metallicity Z is also an important parameter used in seismic modelling, directly affecting the seismic age. A higher value of Z results in a higher seismic mass M . Consequently, the seismic age t of the star appears to be lower, illustration can be seen in ($\alpha_{ov} = 0.2$) in the left panel of Fig. 6. The tracks computed for $Z = 0.008$ and 0.014 are given with dotted and solid lines, respectively.

It demonstrates that an increase of Z results in a shift of the evolutionary tracks towards the lower values of both the luminosity L and effective temperature T_{eff} , resulting in a higher value of the seismic mass M .

Atomic diffusion is responsible for the emergence of irregularities in the horizontal and vertical distribution of chemical elements in stellar atmospheres. It distorts the observed metallicity and hence complicates asteroseismic analyses. Membership of an open cluster can potentially resolve the global metallicity problem, but HD 118660 is a field star located in the solar vicinity at a distance of ~ 70 pc (calculated from the *Gaia* DR3 parallax). Along with that, the estimated iron abundance of the star is close to the solar value (Joshi et al. 2017). In this analysis, we constrain the global metallicity (Z) value close to solar ($Z_\odot = 0.014$; Asplund, Amarsi & Grevesse 2021; Buldgen et al. 2024). Irrespective of the exact value

of Z , HD 118660 is situated within the main-sequence band of the H-R diagram.

3.3.3 Age, luminosity, and effective temperature

Joshi et al. (2017) showed that HD 118660 is a main sequence star. We used this fact to restrict our grid of models within the main-sequence region of the H-R diagram.

The standard luminosity relation is used to calculate an updated value of the luminosity $\log(L/L_\odot)$ of HD 118660. Indeed, a more accurate value of $\pi = 14.2029 \pm 0.0275$ mas for the parallax of HD 118660 has been published recently in the third data release of *Gaia* (Gaia Collaboration 2023). For the interstellar extinction, we used $A_V = 0$ mag. It was derived from the interstellar dust map catalogued by Green et al. (2019). Like Joshi et al. (2017), the bolometric correction $BC = 0.033$ was taken from Flower (1996) and the bolometric magnitude of the sun $M_{\text{bol},\odot} = 4.74$ was published by Willmer (2018), m_v is taken as 6.44 (Gaia Collaboration), which is similar to the mean photometric magnitude observed in *TESS* band in both the sectors. In this way, we derived $\log(L/L_\odot) = 1.09 \pm 0.01$ dex for HD 118660. This value is slightly lower than 1.12 ± 0.27 published by Joshi et al. (2017), but they are compatible if the errors are considered. We adopted the effective temperature $T_{\text{eff}} = 7550 \pm 150$ K measured by Joshi et al. (2017) in combination with our updated value of the luminosity $\log(L/L_\odot) = 1.09 \pm 0.01$ dex to define the observed location of HD 118660 in the H-R diagram in Figs 6 and 7.

One can notice from Fig. 7 that the application of the seismic χ^2 criterion results in stellar models scattered over a large range of T_{eff} and $\log L/L_\odot$. The models at a large distance for the observed




Figure 7. The grey dots show the models with the lowest χ^2 value from each of the 570 evolution tracks mentioned in Section 3.3 overlaid with evolution tracks for $Z = 0.014$ and $\alpha_{ov} = 0.2$. The black error bars show the location of HD 118660 in the H-R diagram.

position of HD 118660 can be safely ignored. At the same time, the strict seismic constraints arising from the characteristics of the identified radial overtones define a dense band that includes the observed position of HD 118660 in the H-R diagram.

3.3.4 Stellar rotation

For the current study, we ignored the effects of stellar rotation in our grid of models. Most CP stars are found to be slow rotators (Conti 1965; Abt & Morrell 1995). However, a recent study showed that the projected rotational velocity $v \sin i$ for Am stars can reach up to 120 km s^{-1} and shows that they are younger compared to non-CP stars with the same rotation rate (Qin et al. 2021). For HD 118660, the available high-resolution spectra have revealed that it is a relatively fast rotator with a $v \sin i$ value of about $100\text{--}110 \text{ km s}^{-1}$ (Catanzaro & Ripepi 2014; Joshi et al. 2017; Gaia Collaboration 2023). Depending on the value of the inclination angle i , the rotation velocity of HD 118660 can be high enough to make our simplified approach unacceptable. To explore this problem in detail, we have analysed the available photometric data for any signature of stellar rotation.

The Keplerian critical (breakup) rotational velocity (Gagnier et al. 2019) can be calculated as

$$v_c = \sqrt{GM/R}, \quad (4)$$

where G is the gravitational constant. By using the values $R = 2.07 R_\odot$ and $M = 1.8 M_\odot$ for the equatorial radius and mass, respectively (second line of Table 2), we find $v_c = 407 \text{ km s}^{-1}$ for HD 118660.

The equatorial rotational velocity v_{eq} is given by

$$v_{eq} = 50.6 R f_{rot}. \quad (5)$$

By using $R = 2.07 R_\odot$ (see Table 2) and by considering $v_c = 407 \text{ km s}^{-1}$ as the maximum and $v \sin i = 108 \text{ km s}^{-1}$ from Joshi et al. (2017) with $i = 90^\circ$ as the minimum value for v_{eq} , respectively, we can estimate the expected range for the rotational frequency f_{rot} of HD 118660 to be $\approx 0.9\text{--}4 \text{ d}^{-1}$.

In the frequency spectra calculated for the *TESS* light curves, we find only two peaks with frequencies falling in the specified range: $f_{23,15} \approx 1.08 \text{ d}^{-1}$ for S23 and $f_{50,16} \approx 1.09 \text{ d}^{-1}$ for S50 (Table 1). If we interpret the average value $1.085 \pm 0.003 \text{ d}^{-1}$ (error

is corresponding to the σ_f of $f_{23,15}$ and $f_{50,16}$ from Table 1) as f_{rot} , then we can use equation (5) to estimate the angle i between the rotational axis and the line of sight. In this way, we derived that $v_{eq} \approx 114 \text{ km s}^{-1}$ and $i \approx 71^\circ \pm 2^\circ$ ($v \sin i$ taken from Joshi et al. 2017). Hence, we find that $v_{eq}/v_c \approx 0.28$. As this value is below the limit of $v_{eq}/v_c = 0.4$ that is set as the limit for the rotational effects to become significant (e.g. Ekström et al. 2012), this would imply that we can consider our modelling of the stellar pulsations of HD 118660 as adequate.

4 CONCLUSIONS

This paper presents a detailed seismic study of HD 118660, a δ Sct pulsator with mild anomalies of its surface chemical composition. Our interest in this particular star is based on its brightness and the accessibility of high-quality and time-resolved photometric and spectroscopic observations. Catanzaro & Ripepi (2014) and Joshi et al. (2017) classify this star as a fast rotator with a $v \sin i$ -value around 100 km s^{-1} , which is close to the observed upper limit for Am stars. Rapid axial rotation remains challenging to account for in asteroseismic modelling and new studies may help to verify the progress achieved in theory. In our study, the effects of rotation are not incorporated. Instead, we are focused on the seismic determination of the mass, radius, and age of HD 118660, as well as on the study of the effects of convective overshooting.

This part of the study is based solely on the currently available *TESS* observations from sectors 23 and 50 and the interpretation and modelling of the observed variations. We estimated the seismic parameters after identifying the pulsation modes detected in the *TESS* photometry. To identify the modes, we used an Échelle diagram and compared the frequency ratios in the theoretical Petersen diagram. This allowed us to conclude that the radial overtones with orders $n = 3 - 6$ are present in the data of HD 118660. Subsequent tests based on the seismic- χ^2 distribution have been used to find the best-fitting model from an appropriate grid of theoretical models calculated with the code CLÉS.

HD 118660 is a late A-type δ Sct pulsator with $T_{eff} = 7550 \pm 150 \text{ K}$ (Joshi et al. 2017) located near to the red edge of the δ Sct instability strip (Dupret et al. 2004). Theory predicts the excitation of high overtones near the blue edge of the instability strip. The observed pulsation pattern of HD 118660 does not seem consistent with these predictions for stars with a solar metallicity. As the global metallicity of HD 118660 hardly differs from the one of the Sun, one can expect that the process of the chemical transport, which is responsible for the chemical peculiarity, modifies the opacity and thus the κ -mechanism in such stars. We intend to address this problem in a follow-up study.

In the current study, we found that the constraints of the radial overtones are not sufficient to constrain the overshooting parameter (α_{ov}) and metallicity (Z). Therefore they are subjected to additional constraints to estimate the seismic parameter. Fig. 7 demonstrates that the known radial overtones can only produce a cloud of luminosities and effective temperatures spread over a huge range. For the final solution, we have selected those stellar models having $\log(L/L_\odot)$ and $\log T_{eff}$ within 1σ from the values derived from observations. The problem with an unconstrained metallicity appears even more critical. In this case, we have adopted $Z = 0.014 \pm 0.002$, based on an *a priori* knowledge of the proximity of the studied star to the Sun. With the metallicity being fixed, the estimated seismic age of HD 118660 ranges within $1.15\text{--}1.35 \text{ Gyr}$ that manifests the star HD 118660 is a ZAMS phase of evolution.

At the same time, we cannot exclude that the real value of the global metallicity of HD 118660 differs slightly (but enough to impact the modelling), depending on the specific origin. We have not elaborated further on this problem as it extends far beyond the scope of the present study. We conclude that our study has shown that with the identified radial modes only, asteroseismology cannot provide sufficient constraints on the convective overshooting and metallicity in δ Sct stars.

Finally, the seismic study of HD 118660 has also confirmed that the star is a fast rotator. In the *TESS* data, we have identified at least one frequency that can most likely be attributed to rotational modulation. This frequency appears in the frequency spectra of *TESS* data obtained in both sectors (S23 and S50). If we interpret it as the rotational frequency f_{rot} and use it in combination with the value for the equatorial stellar radius as found in our study, we could estimate the equatorial rotational velocity to be $\sim 114 \text{ km s}^{-1}$. This value corresponds approximately to 28 per cent of the breakup rotational velocity for an A-type star with the characteristics of HD 118660. We plan to assess the rotational properties of this star in an upcoming paper containing an in-depth spectroscopic study of HD 118660.

ACKNOWLEDGEMENTS

The authors are thankful to the anonymous reviewer for providing insightful comments that significantly improved the manuscript substantially. For this work, MS acknowledges Dr. Simon J. Murphy for helping with the ideas. The work presented here is supported by the Belgo-Indian Network for Astronomy and Astrophysics (BINA), sanctioned by the Department of Science and Technology (DST, Government of India; DST/INT/Belg/P-09/2017) and the Belgian Federal Science Policy Office (BELSPO, Government of Belgium; BL/33/IN12). MS thanks CSIR for supporting the fellowship (09/948(0006)/2020-EMR-1). SJ and OT acknowledge the financial support received from the BRICS grant DST/ICD/BRICS/Call-5/SAPTARISI/2023(G). IY is grateful to the Russian Science Foundation (RSCF grant 21-12-00147) for the partial financial support. AG, DM, and SJ are grateful for the support received from the Indo-Thailand Programme of Co-operation in Science and Technology through the Indo-Thai joint project DST/INT/Thai/P-16/2019. OT acknowledges the International Science Programme (ISP) of Uppsala University for financial support. The PYTHON modules were used in our analysis are LIGHTKURVE, ASTROPY, MATPLOTLIB, SCIPY, and NUMPY.

DATA AVAILABILITY

TESS time-series flux can be found on NASA's MAST website.

REFERENCES

Abt H. A., 2009, *AJ*, 138, 28
 Abt H. A., Morrell N. I., 1995, *ApJS*, 99, 135
 Aerts C., Christensen-Dalsgaard J., Kurtz D. W., 2010, in *Astronomy and Astrophysics Library, Asteroseismology*. Springer Science+Business Media B.V., Germany
 Antoci V. et al., 2014, *ApJ*, 796, 118
 Ashoka B. N. et al., 2000, *Bull. Astron. Soc. India*, 28, 251
 Asplund M., Amarsi A. M., Grevesse N., 2021, *A&A*, 653, A141
 Balona L. A., 2011, *MNRAS*, 415, 1691
 Balona L. A., 2014, *MNRAS*, 437, 1476
 Balona L. A., 2019, *MNRAS*, 490, 2112
 Baran A. S., Koen C., Pokrzywka B., 2015, *MNRAS*, 448, L16

Bedding T. R. et al., 2020, *Nature*, 581, 147
 Breger M., 1979, *PASP*, 91, 5
 Buldgen G., Noels A., Baturin V. A., Oreshina A. V., Ayukov S. V., Scuflaire R., Amarsi A. M., Grevesse N., 2024, *A&A*, 681, A57
 Catanzaro G., Ripepi V., 2014, *MNRAS*, 441, 1669
 Christensen-Dalsgaard J., Arentoft T., Brown T. M., Gilliland R. L., Kjeldsen H., Borucki W. J., Koch D., 2008, *Commun. Asteroseismol.*, 157, 266
 Claret A., Torres G., 2016, *A&A*, 592, A15
 Claret A., Torres G., 2017, *ApJ*, 849, 18
 Claret A., Torres G., 2018, *ApJ*, 859, 100
 Claret A., Torres G., 2019, *ApJ*, 876, 134
 Constantino T., Baraffe I., 2018, *A&A*, 618, A177
 Conti P. S., 1965, *ApJ*, 142, 1594
 Deal M., Monier R., 2020, *AAS*, 4, 144
 Donati J. F., Landstreet J. D., 2009, *ARA&A*, 47, 333
 Dornan V., Lovekin C. C., 2022, *ApJ*, 924, 130
 Dupret M. A., Grigahcène A., Garrido R., Gabriel M., Scuflaire R., 2004, *A&A*, 414, L17
 Dupret M. A., Grigahcène A., Garrido R., Gabriel M., Scuflaire R., 2005, *A&A*, 435, 927
 Ekström S. et al., 2012, *A&A*, 537, A146
 Fleming S. W., 2019, *American Astronomical Society Meeting Abstracts* #233, p. 245.11
 Flower P. J., 1996, *ApJ*, 469, 355
 Fossati L., Bagnulo S., Monier R., Khan S. A., Kochukhov O., Landstreet J., Wade G., Weiss W., 2007, *A&A*, 476, 911
 Fossati L., Bagnulo S., Landstreet J., Wade G., Kochukhov O., Monier R., Weiss W., Gebran M., 2008, *A&A*, 483, 891
 Gagnier D., Rieutord M., Charbonnel C., Putigny B., Espinosa Lara F., 2019, *A&A*, 625, A88
 Gaia Collaboration, 2023, *A&A*, 674, A1
 Goupil M. J., Dupret M. A., Samadi R., Boehm T., Alecian E., Suarez J. C., Lebreton Y., Catala C., 2005, *JA&A*, 26, 249
 Green G. M., Schlafly E., Zucker C., Speagle J. S., Finkbeiner D., 2019, *ApJ*, 887, 93
 Guzik J. A., 2021, *Front. Astron. Space Sci.*, 8, 55
 Guzik J. A., Fryer C., Fontes C. J., 2018, in Wade G. A., Baade D., Guzik J. A., Smolec R. eds, 3rd BRITE Sci. Conf., Opacity Effects on Pulsations of A-Type Stars, Vol. 8, p. 59
 Guzik J. A., Garcia J. A., Jackiewicz J., 2019, *Front. Astron. Space Sci.*, 6, 40
 Hansen C. J., Cox J. P., van Horn H. M., 1977, *ApJ*, 217, 151
 Hey D., Aerts C., 2024, *A&A*, 688, A93
 Hey D., Ball W., 2022, *Astrophysics Source Code Library*, record: 2207.005
 Holdsworth D. L. et al., 2021, *MNRAS*, 506, 1073
 Holdsworth D. L. et al., 2024, *MNRAS*, 527, 9548
 Houdek G., 2000, in Breger M., Montgomery M. eds, *ASP Conf. Ser.*, Vol. 210, Delta Scuti and Related Stars. Astron. Soc. Pac., San Francisco, p. 454
 Howell S. B. et al., 2014, *PASP*, 126, 398
 Hubrig S., Järvinen S. P., Alvarado-Gómez J. D., Ilyin I., Schöller M., 2023, *MNRAS*, 526, L83
 Johnston C., 2021, *A&A*, 655, A29
 Joshi S. et al., 2003, *MNRAS*, 344, 431
 Joshi S., Mary D. L., Martinez P., Kurtz D. W., Girish V., Seetha S., Sagar R., Ashoka B. N., 2006, *A&A*, 455, 303
 Joshi S., Mary D. L., Chakradhari N. K., Tiwari S. K., Billaud C., 2009, *A&A*, 507, 1763
 Joshi S., Ryabchikova T., Kochukhov O., Sachkov M., Tiwari S. K., Chakradhari N. K., Piskunov N., 2010, *MNRAS*, 401, 1299
 Joshi S. et al., 2012, *MNRAS*, 424, 2002
 Joshi S. et al., 2016, *A&A*, 590, A116
 Joshi S., Semenko E., Moiseeva A., Sharma K., Joshi Y. C., Sachkov M., Singh H. P., Yerra B. K., 2017, *MNRAS*, 467, 633
 Joshi S. et al., 2022, *MNRAS*, 510, 5854

- Kjeldsen H., Bedding T. R., 2012, in Griffin E., Hanisch R., Seaman R.eds, *New Horizons in Time-Domain Astronomy*, Proceedings of the International Astronomical Union, IAU Symposium, Vol. 285,p. 17
- Kurtz D. W., 1982, *MNRAS*, 200, 807
- Kurtz D. W., 2022, *ARA&A*, 60, 31
- Kurtz D. W., Hubrig S., González J. F., van Wyk F., Martinez P., 2008, *MNRAS*, 386, 1750
- Ledoux P., 1949, *Mem. Soc. R. Sci. Liege*, 9, 3
- Lee Y.-H., Kim S. S., Shin J., Lee J., Jin H., 2008, *PASJ*, 60, 551
- Lenz P., Breger M., 2005, *Commun. Astroseismol.*, 146, 53
- Lightkurve Collaboration, 2018, *Astrophysics Source Code Library*, record ascl soft:1812.013
- Lovekin C. C., Guzik J. A., 2017, *ApJ*, 849, 38
- Martinez P. et al., 2001, *A&A*, 371, 1048
- Michaud G., 1970, *ApJ*, 160, 641
- Michaud G., Alecian G., Richer J., 2015, *Astronomy and Astrophysics Library*, Atomic Diffusion in Stars. Springer International Publishing, Switzerland
- Montgomery M. H., O'Donoghue D., 1999, *Delta Scuti Star Newsl.*, 13, 28
- Morgan D., Richard M., 2020, *AAS*, 4, 144, Res. Notes
- Murphy S. J., 2014, PhD thesis, Univ. Central Lancashire
- Murphy S. J., Hey D., Van Reeth T., Bedding T. R., 2019, *MNRAS*, 485, 2380
- Murphy S., Gray R., Corbally C., Kuehn C., Bedding T., Killam J., 2020, *MNRAS*, 499, 2701
- Murphy S. J., Saio H., Takada-Hidai M., Kurtz D. W., Shibahashi H., Takata M., Hey D. R., 2020, *MNRAS*, 498, 4272
- Murphy S. J., Joyce M., Bedding T. R., White T. R., Kama M., 2021, *MNRAS*, 502, 1633
- Neiner C., Lampens P., 2015, *MNRAS*, 454, L86
- Petersen J. O., 1973, *A&A*, 27, 89
- Petersen J. O., 1978, *A&A*, 62, 205
- Preston G. W., 1974, *ARA&A*, 12, 257
- Qin L. et al., 2021, *AJ*, 162, 32
- Reese D. R., 2022, *Front. Astron. Space Sci.*, 9, 934579
- Renson P., Manfroid J., 2009, *A&A*, 498, 961
- Ricker G. R. et al., 2015, *J. Astron. Telesc. Instrum. Syst.*, 1, 014003
- Rodríguez E., Balona L. A., López-González M. J., Ocampo S., Martín-Ruiz S., Rodríguez-López C., 2020, *MNRAS*, 491, 4345
- Sarkar M., Joshi S., de Cat P., 2024, *Bull. Soc. R. Sci. Liège*, 93, 285
- Scuflaire R., Théado S., Montalbán J., Miglio A., Bourge P. O., Godart M., Thoul A., Noels A., 2008a, *Ap&SS*, 316, 83
- Scuflaire R., Montalbán J., Théado S., Bourge P. O., Miglio A., Godart M., Thoul A., Noels A., 2008b, *Ap&SS*, 316, 149
- Sikora J. et al., 2019, *MNRAS*, 487, 4695
- Sikora J., Wade G. A., Rowe J., 2020, *MNRAS*, 498, 2456
- Solano E., Fernley J., 1997, *A&AS*, 122, 131
- Tassoul M., 1980, *ApJS*, 43, 469
- Trust O., Jurua E., De Cat P., Joshi S., 2020, *MNRAS*, 492, 3143
- Trust O., Jurua E., De Cat P., Joshi S., Lampens P., 2021, *MNRAS*, 504, 5528
- Trust O., Mashonkina L., Jurua E., De Cat P., Tsymbal V., Joshi S., 2023, *MNRAS*, 524, 1044
- Umezu M., 1995, *MNRAS*, 276, 1287
- Uytterhoeven K. et al., 2011, *A&A*, 534,A125
- Willmer C. N. A., 2018, *ApJS*, 236, 47

This paper has been typeset from a $\mathrm{T}_{\mathrm{E}}\mathrm{X}/\mathrm{L}^{\mathrm{A}}\mathrm{T}_{\mathrm{E}}\mathrm{X}$ file prepared by the author.