




Binary stars in the space era
Keele University – Friday 4th July 2025




Exploring spectroscopic binaries: From SB9 to the 4MOST large survey

Thibault Merle
Université Libre de Bruxelles
Observatoire Royal de Belgique




Why spectroscopic binaries (SB)?

Inensitive to the distance



Wide range of orbital periods



- The SB9 catalogue and its evolution
 - History
 - Content including triples and quadruples
 - Evolution
- Comparison of SB9 with Gaia DR3
 - Gaia color-absolute magnitude diagram
 - SB9 x NSS common sample
 - Its characteristics
- Spectroscopic binaries in 4MOST
 - A massive MOS survey
 - Many SB projects
 - Multiple stars detection and characterisation




History of the **SB9** catalogue

v6 – v8




Dominion Astrophysical Observatory
Batten (1967) 740 SB

v1 – v5



Lick Observatory
Campbell & Curtis (1905) 140 SB

Thibault Merle



<https://sb9.astro.ulb.ac.be>



Binary Stars in the Space Era



2000: IAU initiative
from commission G1



ULB

SB9
Pourbaix+ (2004)
2390 SB

↓
2021
4000 SB

- 5000 orbits
 - 4000 systems
- 70% SB1
30% SB2


Content of **SB9** catalogue



Content of **SB9** catalogue

Hierarchies

- ~75 spectroscopic triples
- ~30 spectroscopic quadruples (including 3 optical)
- 3 quadruples (2+2 configuration) with complete orbits:
 - μ ori
 - HD 117078
 - VW LMi
- 5 systems with multiplicity > 4 having at least 2 subsystems in SB9:
 - 3 quintuplets (1 in the Orion Trapezium),
 - and 2 sextuplets (3+3 and 4+2 configurations)






The eXtended catalogue of SB orbits


<http://astro.ulb.ac.be/sbx>

Poster 16

- **SB9** is outdated in web design, data access and storage
- **SBX** has a PostgreSQL database management system
- **SBX** uses a Simbad name resolver API
- **SBX** is following IVOA standards with a TAP service using ADQL



Gaia color-magnitude diagram of SB9 systems





Not in Gaia DR3:

- The 21 brightest systems
- 6 faint systems
- 18 duplicates

Late type SB are under-represented

Cataclysmic variables
MS+WD binaries


Gaia color-magnitude diagram of SB9 systems




SB9 x Gaia DR3 NSS common sample

828 binaries:

- This only represent 1/5 of SB9 catalogue
- Many SB9 binaries with periods larger than 1000 d
- Some Gaia binaries with high eccentricities at short periods




SB9 x Gaia DR3 NSS common sample



P_{Gaia} VS. P_{SB9}

Selection 1: $|\Delta P|/P < 0.1$
 $\Rightarrow \sim 70$ outliers



e_{Gaia} VS. e_{SB9}

Selection 2: $|\Delta e| < 0.1$
 $\Rightarrow \sim 170$ outliers


Clean sample
= selection 1 & selection 2

Common sample: 850
Clean sample: 660

SB9 x Gaia DR3 NSS common sample

Common sample of 828 binaries:


- This only represent 1/5 of SB9 catalogue
- Many SB9 binaries with periods larger than 1000 d
- Some Gaia binaries with high eccentricities at short periods



SB9 x Gaia DR3 NSS common sample

Common sample of 828 binaries:


- This only represent 1/5 of SB9 catalogue
- Many SB9 binaries with periods larger than 1000 d
- Some Gaia binaries with high eccentricities at short periods






SB9 x Gaia DR3 NSS common sample

Common sample of 828 binaries:

- This only represent 1/5 of SB9 catalogue
- Many SB9 binaries with periods larger than 1000 d
- Some Gaia binaries with high eccentricities at short periods



SB9 x Gaia DR3 NSS common sample



Merle+, in prep.

- Only 20% of SB9 systems in NSS.
- Increase to 25% when restricting to the same period and magnitude ranges.
- Many internal filtering processes in DR3 (Gosset+ 2025, Damerджи, in prep.)

4
MOST



A massive spectroscopic survey






- 4-m Multi-Object Spectroscopic Telescope on VISTA/ESO
- 2400 fibres per single exposure
- 4 square degrees field of view
- Optical wavelength coverage
- Low-resolution: 4 000 – 8 000, 1 600 fibres, $V_{\max} \sim 20$
- High resolution: $\sim 20\,000$, 800 fibres, $V_{\max} \sim 16$
- 5 y survey starting in 2026



A massive spectroscopic survey

- 10 consortium surveys: 70%
- 14 ESO community surveys: 30%
- 9 infrastructure working groups (IWG) including a Stellar Multiplicity Working Group as part of IWG2 (survey strategy and simulations)
- Implementation of cadence:
 - For free
 - On request

S0401 - 4MIDABLEHR_DISK, N targets: 6090427



<https://www.4most.eu/cms/science/overview/>



Spectroscopic binaries in 4MOST

Binary nature of SB candidates (LR/HR) - T. Merle

Dependence of binary fraction on metallicity (LR/HR) - G. Traven

Composite-colour hot subdwarf binaries - M. Dorsch

Detecting spectroscopic stellar systems – T. Merle

MULTIFIT pipeline and parameter biases (LR/HR) – M. Dorsch

Population properties of binary stars in our Galaxy – G. Traven, **B. Jamnik (see poster P41)**

Double white dwarf binaries – **I. Pelisoli (talk on Tuesday)**

WDs with unseen low-mass MS (or brown dwarf) companions – Rebassa-Mansergas

Dormant Black Holes survey overview and catalogue – M. Pawlak

Abundance analysis of multiple stars – H. Ernandes

Spectroscopic binarity of the Gaia astrometric binaries – T. Mazeh

The distribution of the rotational broadening of all 4MOST spectra – T. Mazeh

Characterising sources with significant astrometric excess noise (LR/HR) – P. Gandhi

Fundamental parameters (mass, radius) of eclipsing binaries – T. Merle

Wolf-Rayet stars and spectral disentangling of SB1 – **T. Shenar**

Etc.



4

MOST

Multiple stars detection & characterization

Composite spectrum =
Main sequence F + subgiant K

MULTIFIT from M. Dorsch
Approach similar to El-Badry+ (2018) in APOGEE
Using neural networks (The Payne, Ting+ 2017)



$$\chi^2_{\text{SB}} = \chi^2_{\text{spec}} + \chi^2_{\text{phot}}$$

4 MOST

Binary nature of many candidates

For unresolved SB2 (~15%):

- Derivation of atmospheric parameters

For SB1 (~1.5%):

- Minimum RV dispersion for SB1 detection
- Set upper limits on orbital periods
- Estimation of the period distribution


For SB2 (few thousands):

- Partial orbital characterization
- Derivation of atmospheric parameters

For <1% of SB_n, n≥1:

- Full orbital characterization
- Full atmospheric characterization

In addition few tens to hundreds of SB2 will be turn into triples or quadruples.





- **SB9 Catalogue and Its Evolution**

The SB9 catalogue compiles thousands of spectroscopic binary orbits, including higher-order multiples, and has evolved into the modern SBX database with improved data access and standards.

<http://astro.ulb.ac.be/sbx>



- **Comparison with Gaia DR3**

Cross-matching SB9 with Gaia DR3 NSS reveals only a partial overlap (about 20–25%), highlighting differences in detection limits, period coverage, and the need for further investigation into the completeness of binary catalogues.

- **Spectroscopic Binaries in the 4MOST Survey**

The upcoming 4MOST survey will massively expand spectroscopic binary detection, enabling detailed studies of binary fractions, period distributions, and higher-order multiples across the Galaxy, with new tools and strategies for characterizing both resolved and unresolved systems.

Gilmore girls





- **SB9 Catalogue and Its Evolution**

The SB9 catalogue compiles thousands of spectroscopic binary orbits, including higher-order multiples, and has evolved into the modern SBX database with improved data access and standards.

<http://astro.ulb.ac.be/sbx>



- **Comparison with Gaia DR3**

Cross-matching SB9 with Gaia DR3 NSS reveals only a partial overlap (about 20–25%), highlighting differences in detection limits, period coverage, and the need for further investigation into the completeness of binary catalogues.



- **Spectroscopic Binaries in the 4MOST Survey**

The upcoming 4MOST survey will massively expand spectroscopic binary detection, enabling detailed studies of binary fractions, period distributions, and higher-order multiples across the Galaxy, with new tools and strategies for characterizing both resolved and unresolved systems.

RV curves: SB9 vs Gaia DR3 NSS

